

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 30-09-2014		2. REPORT TYPE Final		3. DATES COVERED (From - To) 01-JAN-2011 to 30-SEP-2014	
4. TITLE AND SUBTITLE Seabed Geoacoustic Structure at the Meso-Scale				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER N00014-11-1-0124	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Charles W. Holland, Ph.D.				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Pennsylvania State University Applied Research Laboratory P. O. Box 30 State College, PA 16804-0030				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of naval Research 875 North Randolph Street Arlington, VA 22203-1995				10. SPONSOR/MONITOR'S ACRONYM(S) ONR	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					

14. ABSTRACT

The long term science goals of the project are to understand the nature of the seabed at the geoacoustic meso-scale $O(10^0-10^3)$ m and determine how these structures impact acoustic propagation, diffuse reverberation, and clutter. Several discoveries were made in the course of the project, notably that a) there is a persistent scale, $O(0.1)$ m that controls reverberation in a number of shallow water environments, and b) clutter can be generated by non-discrete, i.e., smoothly and slowly-varying sediment structures (in addition to traditional discrete features, e.g., wrecks). In addition, there were a number of significant advances including: development of measurement techniques/processing to measure meso-scale geoacoustic variability over tens of km with resolution of 0.1 m/7 m in the vertical/lateral dimension; and developing tools to separate frequency-dependent effects so that unbiased estimates of intrinsic attenuation can be made. Even on the mid to outer shelf, substantial lateral variability at the 10 m lateral scale is observed. The underlying processes that lead to that strong variability are not understood at this time.

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)	
U	U	U	U	4		

20150406 011

Seabed Geoacoustic Structure at the Meso-Scale

Charles W. Holland

The Pennsylvania State University
Applied Research Laboratory
P.O. Box 30, State College, PA 16804-0030
Phone: (814) 865-1724 Fax (814) 863-8783 email: holland-cw@psu.edu

Grant Number: N00014-11-1-0124

LONG TERM GOALS

The long term science goals are to understand the nature of the seabed at the geoacoustic meso-scale $O(10^0-10^3)$ m and determine how these structures impact acoustic propagation, diffuse reverberation, and clutter.

OBJECTIVES

The objectives are to develop new observational methods to quantify meso-scale seabed variability/uncertainty and also develop modeling techniques to understand the impact of spatial variability on propagation and reverberation.

RESULTS

The scientific results are contained in 15 peer-reviewed publications. Some of the highlights are briefly described below.

Discoveries

- discovery of a persistent scale that controls reverberation in a number of shallow water environments (Ref[12]). The consistency of the scale, $O(0.1)$ m, suggests an underlying feature or mechanism (as yet unknown) that is consistent across many ostensibly diverse geological settings. This has significant implications for employment of active sonar, and also for seabed database development.
- Discovered that clutter can be generated by non-discrete, i.e., smoothly and slowly-varying sediment structures (Ref [13]). Generally, clutter from the seabed has been associated with discrete features, e.g., wrecks or carbonate mounds. This discovery has implications for sonar systems and for acoustic probing of sediment structures. (with Dale Ellis, DRDC-A)

Significant Advances

- Developed new method (Ref [9]) for estimating dispersion that explicitly includes separating its frequency-dependent effects from the frequency-dependent effects of sediment layering and gradients. The method also has the virtue of determining the attenuation mechanism, i.e.,

friction or viscous losses (which itself can be a function of frequency). (with Jan Dettmer, U. Victoria)

- Measurements of 1000-3600 Hz attenuation in a muddy sediment fabric using long-range diffuse reverberation (Ref [6]). The importance of this is that
 - a) there is a glaring paucity of measurements in the 600 – 4000 Hz band
 - b) mud attenuation has historically been very difficult to measure at all frequencies due to measurement biases on high frequency samples
 - c) uncertainties in the measurements are almost never reported. Our measurements showed quite low uncertainties – average attenuation in a 10 m thick mud layer was 0.009 ± 0.003 dB/m/kHz. (with Stan Dosso, U. Victoria)
- In collaboration with Gavin Steininger and others at U. Victoria, developed models and methods, Ref [2] to :
 - a) determine whether seabed scattering was dominated by interface roughness or sediment volume heterogeneities;
 - b) determine the probability of volume heterogeneities as a function of depth in the sediment stack; and
 - c) determine the outer scale of the heterogeneities.

The ability to determine the scattering mechanism (e.g., interface roughness versus volume heterogeneities) from acoustic data is important for both scientific and operational purposes. For the former, it will permit a deeper understanding of the seabed processes that lead to scattering; for the latter, an emerging bottom scattering database will require provincing methods (e.g., is it possible to extrapolate observations in one area to another that has no measurements?). This, and other provincing questions are greatly aided by identification of the dominant scattering mechanism.

- Developed two distinct techniques to measure/analyze meso-scale seabed variability in shallow water using acoustic reflection data from an AUV. One technique (Ref [15]), exploits the high information content of the reflection coefficient versus angle and frequency, estimating all the geoacoustic parameters and their uncertainties with depth and along the track, i.e., a 2D model with a lateral resolution of about 7 m (with Peter Nielsen, CMRE, Jan Dettmer, U. Victoria). The other technique is several orders of magnitude faster computationally and estimates the 2D sound speed profile using the image source method with a lateral resolution of ~50 m. See Ref [4],[8]). (with Samuel Pinson, post-doc).

Incremental advances

- Developed technique to measure low-angle bi-static seabed scattering in shallow water using an AUV (Ref [16]). (with Peter Nielsen, CMRE, and Chad Smith, PSU)
- In collaboration with Gavin Steininger, Stan Dosso and Jan Dettmer (all from U. Vic) estimated roughness parameters from direct path seabed backscatter data (Refs [10],[11]). This represents the first time there has been an objective inverse method applied to seabed acoustic scattering data to provide not only the scattering parameters, but also parameter uncertainties.
- Conducted *in-situ* geoacoustic measurements of sediments possessing a very wide range of grain sizes including very large granules (e.g., shells, pebbles and cobbles), see Ref [1]. This is a sediment type for which no sediment acoustic models currently exist, and very little has been published. However, this kind of sediment may in fact be relatively common. The results are intended to provide geoacoustic data where little exists in the literature and also to motivate development of models that can treat a wide range of grain sizes. (with partners at U. Victoria).

RELATED PROJECTS

ONR shallow water field experiments: the advances in the project described above will motivate and inform experiment design to disentangle effects of sediment dispersion from other frequency dependent effects.

SPAWAR PMW-120 Seabed Bottom Scattering database development. Many aspects of the theoretical and measurements-related research under the ONR research described above have informed this development.

PUBLICATIONS

- [1] Holland C.W., J.D. Dettmer, G. Steininger, S.E. Dosso, A. Lowrie, Acoustic measurements of sediments with pebbles and cobbles, *J. Acoust. Soc. Am.*, in review, 2014.
- [2] Steininger G., S. Dosso, C.W. Holland and J. Dettmer, Estimating seabed scattering mechanisms via Bayesian model selection, *J. Acoust. Soc. Am.*, 136, 1552-1562, 2014.
- [3] Dosso S.D., J. Dettmer, Steininger G., and C.W. Holland, Efficient trans-dimensional Bayesian inversion for geoacoustic profile estimation, *J. Inv. Problems*, 30, 2014, doi:10.1088/0266-5611/30/11/114018,
- [4] Pinson S. and C.W. Holland, Seafloor sound-speed profile and interface dip angle characterization by the image source method, *J. Acoust. Soc. Am.*, 136, 596-603, 2014.
- [5] Steininger G., S. Dosso, C.W. Holland and J. Dettmer, A trans-dimensional polynomial-spline parameterization for gradient-based geoacoustic inversion, *J. Acoust. Soc. Am.*, 136, 1563-1573, 2014.
- [6] Holland C.W. and S.E. Dosso, Mid frequency shallow-water fine-grained sediment attenuation from waveguide reverberation, *J. Acoust. Soc. Am.*, 134, 131-134, 2013.
- [7] Dettmer J, C.W. Holland and S.E. Dosso, Trans-dimensional uncertainty estimation for dispersive seabed sediments, *Geophysics*, 78, WB63-WB76, 2013.
- [8] Pinson S., L. Guillon and C.W. Holland, Range dependent sound speed profile characterization with an horizontal array by the image source method, *J. Acoust. Soc. Am.*, 134, 156-165, 2013.
- [9] Holland C.W. J. Dettmer, In-situ sediment dispersion estimates in the presence of discrete layers and gradients, *J. Acoust. Soc. Am.*, 133, 50-61, 2013.
- [10] Steininger G., C.W. Holland, S. Dosso and J. Dettmer, Seabed roughness parameters from joint backscatter and reflection inversion on the Malta Plateau, *J. Acoust. Soc. Am.*, 134, 1833-1842, 2013.
- [11] Steininger G., J. Dettmer, S. Dosso, and C.W. Holland, Trans-dimensional joint inversion of seabed scattering and reflection data, *J. Acoust. Soc. Am.*, 133, 1347-1357, 2013.
- [12] Holland C.W., Evidence for a common scale $O(0.1)$ m that controls seabed scattering and reverberation in shallow water, *J. Acoust. Soc. Am.*, 132, 2232-2238, 2012.

- [13] Holland, C.W. and D.D. Ellis, Clutter from non-discrete seabed structures, *J. Acoust. Soc. Am.*, 131, 4442-4449, 2012.
- [14] Dosso S. E., C.W. Holland, M. Sambridge, Parallel tempering for strong nonlinear problems, *J. Acoust. Soc. Am.*, 132, 3030-3040, 2012.
- [15] Holland C.W., P.L. Nielsen, J. Dettmer, and S.E. Dosso, Resolving meso-scale seabed variability using reflection measurements from an autonomous underwater vehicle, *J. Acoust. Soc. Am.*, 131, 1066-1078, 2012.
- [16] Holland, C.W., C.M. Smith, and P.L. Nielsen, Bistatic seabed scattering measurements from an autonomous undersea vehicle, *European Conference on Underwater Acoustics*, Edinburg, UK, 2012.
- [17] Guillon L., Holland C.W. and C. Barber, Cross-spectral analysis of low-frequency acoustic waves reflected by the seafloor, *IEEE J. Ocean Eng.*, 36, 248-258, 2011.